Plants small to large, forming dense tufts, dark green, often lustrous. **Stems** erect, simple to few branched; in cross-section central strand present; radiculose, occasionally tomentose **Leaves** equally distant or crowded distally, crispate to ± flexuose and erect to erect-spreading when dry, erect-spreading to spreading when wet, lanceolate, oblong-lanceolate, ovate, obovate-oblong, apex short to long acuminate, base slightly decurrent or not; margins limbate, plane above, often reflexed to recurved in lower 1/2, entire to more commonly serrulate or serrate above; costae single, usually strong, short to long excurrent, rarely subpercurrent, awn smooth or toothed; median cells mostly thin-walled, broadly hexagonal or short to long rhomboidal-hexagonal, rhomboidal toward margin and apex; lower and basal cells short rectangular; marginal border cells linear and thick-walled, 1-4 rows. **Gemmae** often present, as axillary bulbous gemmae in some species. **Dioicous. Perichaetia** terminal; leaves often larger. **Setae** elongate, erect to weakly flexuose, slender to stout, smooth. **Capsules** suberect to commonly erect, urn obloid-cylindrical, rarely subglobose, slightly asymmetric, neck usually distinct; opercula conic, rarely rostrate; peristome double, exostome teeth 16, joined at base, distally papillose (fragile), trabeculae well developed; endostome basal membrane low, segments somewhat well developed to rudimentary, narrowly perforate, cilia absent or reduced. **Calyptrae** cucullate, smooth and naked. **Spores** spherical, lightly papillose.
Plantas pequeñas hasta grandes, formando manojos densos, de color verde oscuro, a menudo brillantes. **Tallos** erectos, simples hasta poco ramificados; cordón central presente en corte transversal; radiculosos, ocasionalmente tomentosos. **Hojas** equidistantes o agregadas distalmente, crispas hasta ± flexuosas y erectas hasta erecto-extendidas en seco, erecto-extendidas patentes en húmedo, lanceoladas, oblongo-lanceoladas, ovadas, obovado-oblongas, ápices cortos hasta largo-acuminado, bases escasamente decurrentes o no; márgenes limbados, planos hacia el ápice, a menudo reflexos hasta recurvados en la ½ inferior, enteros a más comúnmente serrulados o serrados hacia el ápice; costa simple, generalmente fuerte, corto- hasta largo-excurrente, rara vez subpercurrente, arista lisa o dentada; células mediales por lo general con paredes delgadas, anchamente hexagonales o corto- hasta largo- romboidal-hexagonales, rómbicas cerca al margen y ápice; células inferiores y basales corto-rectangulares; margen bordeado por células lineales y con paredes gruesas, 1-4 filas. **Yemas** a menudo presentes, como yemas bulbosas axilares en algunas especies. **Dioicas**. **Periquecios** terminales; hojas a menudo mayores. **Setas** alargadas, erectas hasta débilmente flexuosas, delgadas hasta robustas y lisas. **Cápsulas** suberectas hasta comúnmente erectas, urna obloide-cilíndrica, rara vez subglobosa, escasamente asimétrica, cuello generalmente diferenciado; opérculo cónico, rara vez rostrado; peristoma doble, exostoma con 16 dientes, unidos en la base, distalmente papilosos (frágiles), trabéculas bien desarrolladas; membrana basal del endostoma baja, segmentos algo desarrollados hasta rudimentarios, angostamente perforados, cilios ausentes o reducidos. **Caliptra** cuculada, lisa y desnuda. **Esporas** esféricas, finamente papilosas.