Plants small to medium sized, forming loose, thin mats, glossy green, yellowish-green or golden. **Stems** creeping and spreading, irregularly pinnately branched, leafy stems strongly complanate-foliate or erect and arching distally; in cross-section central strand absent; paraphyllia and apparently pseudoparaphyllia absent; rhizoids often clustered below. **Leaves** crispate or not when dry; stem leaves differentiated between dorsal and lateral, **lateral leaves**, ovate, oblong, orbicular or ovate- to oblong-lanceolate, 0.5-2.0 mm long, to 1.2 mm wide, apex acute to short acuminate, often slightly keeled, base subauriculate to rounded; margins usually recurved in part or throughout, doubly-dentate or -serrate, occasionally single serrate between; ecostate; median cells linear to linear-vermicular, firm-walled, smooth; basal cells oblong-fusiform or-rhomboidal, porose; **dorsal leaves**, shorter, usually broader, ovate, oval to orbicular. **Gemmae**, present beneath stems or on specialized branches with differentiated leaves, cylindrical. **Dioicous**. **Perichaetia** lateral, leaves differentiated. **Setae** elongate, 4-12 mm long, papillose throughout or distally. **Capsules** suberect to inclined, urn ovoid to elliptical, 0.5-1.5 mm long, exothecial cells collenchymatous; opercula conic-rostrate; peristome, double, exostome teeth papillose, endostome basal membrane low to high, segments weakly perforate and keeled, spiculose, cilia absent. **Calyptrae** mitrate, base ciliate. **Spores** spherical, finely papillose.

Plantas pequeño hasta medianas, formando matas, laxas y delgadas, verde brillantes, verde-amarillentas o doradas. **Tallos** reptantes y patentes, ramificaciones irregularmente pinnadas, tallos foliosos marcadamente complanado-foliosos o erectos y arqueados distalmente; en corte transversal sin cordón central; parafilos y aparentemente pseudoparafilos ausentes; rizoides a menudo agregados abajo. **Hojas** crispas o no en seco; hojas de los tallos diferenciadas entre dorsales y laterales, hojas laterales, ovadas, oblongas, orbiculares u ovado- hasta oblongo-lanceoladas, 0.5-2.0 mm de largo, hasta 1.2 mm ancho, ápices agudos hasta corto-acuminados, a menudo escasamente carinados, bases subauriculadas hasta redondeadas; márgenes generalmente recurvados parcial o completamente, doble-dentados o -serrados, ocasionalmente intercalados con dientes simples; ecostadas; células mediales lineales hasta lineal-vermiculares, conm paredes firmes, lisas; células basales oblongo-fusiformes o romboidales, porosas; hojas dorsales, cortas, generalmente más anchas, ovadas, ovales hasta orbiculares. **Yemas**, presentes en la porción inferior de los tallos o en ramas especializadas con hojas diferenciadas, cilíndricas. **Dioicas**. **Periquecios** laterales, hojas diferenciadas. **Setas** alargadas, 4-12 mm de largo, completa o distalmente papilosas. **Cápsulas** suberectas hasta inclinadas, urna ovoide hasta elíptica, 0.5-1.5 mm de largo, células exoteciales colenquimatosas; opérculo cónico-rostrado; peristoma, doble, exostoma con dientes papilosos, membrana basal del endostoma baja hasta alta, segmentos débilmente perforados y carinados, espiculadas, cilios ausentes. **Caliptra** mitrada, base ciliada. **Esporas** esféricas, finamente papilosas.