Plants mostly small to somewhat medium sized, usually forming small tufts, occasionally solitary, glossy pale green to golden-brown or yellow. **Primary stems** short creeping, inconspicuous. **Secondary stems** erect or suberect, several branched, spirally foliate. **Leaves** crowded, broadly lanceolate to linear-lanceolate, apex short to long acuminate, rarely cuspidate, base rounded; margins plane or recurved one or both sides, entire or apices rarely serrulate, weakly to strongly limbate; costae single, 1/2-4/5 lamina length; laminal cells smooth, upper and median cells oval to fusiform or linear, thick-walled, smooth; basal and insertion cells linear to oblong or rectangular, often yellowish-brown to golden; marginal cells long linear, forming a distinct border of few to many rows. **Autoicous. Perichaetia** lateral; leaves much smaller than stem leaves, oblong- to ovate-short lanceolate. **Setae** elongate, slender to stout, smooth to more commonly papillose-roughened throughout or distally. **Capsules** erect to suberect, urn ovoid; exothecial cells collenchymatous, or walls equally thickened; stomata on neck or urn base, superficial; opercula conic-short rostrate; peristome double, exostome teeth 16, narrowly lanceolate, papillose, not furrowed; endostome basal membrane low, segments 16, linear, lightly to strongly papillose, keeled and perforate. **Calyptrae** campanulate, base fringed with hairs, distally smooth to somewhat roughened. **Spores** spherical to ovoid, mostly lightly papillose.

Plantas generalmente pequeñas hasta algo medianas, generalmente formando manojos pequeños, ocasionalmente solitarias, de color verde-pálido brillante hasta pardo-dorado o amarillo. **Tallos primarios** corto-reptantes, inconspícuos. **Tallos secundarios** erectos o suberectos, varias veces ramificados, espiralmente foliosos. **Hojas** agregadas, anchamente lanceoladas hasta lineo-lanceoladas, ca. 2-4 mm de largo, ápices cortos hasta largo-acuminados, rara vez cuspidados, base redondeada; márgenes planos o recurvados en uno o ambos lados, ápices enteros o rara vez serrulados, débilmente a marcadamente limbadas; costa simple, ocupando 1/2-4/5 de la longitud de la lámina; células de la lámina lisas, células superiores y mediales ovales hasta fusiformes o lineales, con paredes gruesas, lisas; células basales y de la inserción lineales hasta oblongas o rectangulares, a menudo pardo-amarillentas a doradas; células del margen largo-lineales, formando un borde conspicuo de pocas a muchos filas. **Autoicas**. **Periquecios** laterales; hojas menores que las hojas del tallo, oblongas hasta ovadas corto-lanceoladas. **Setas** alargadas, hasta 7-15 mm de largo, delgadas hasta robustas, lisas a más comúnmente papiloso-rugosas en toda la superficie o en la parte distal. **Cápsulas** erectas hasta suberectas, urna ovoide; células exoteciales colenquimatosas, o paredes igualmente engrosadas; estomas en el cuello o la base de la urna, superficiales; opérculo corto-cónico rostrado; peristoma doble, exostoma con 16 dientes, angostamente lanceolados, papilosos, no acanalados; membrana basal del endostoma baja, con 16 segmentos, lineales, ligera hasta marcadamente papilosos, carinados y perforados. **Caliptra** campanulada, base ciliada, distalmente lisa hasta algo rugosa. **Esporas** esféricas hasta ovoides, por lo general finamente papilosas.