Plants small to occasionally medium sized, forming tufts or thin mats, glossy green to yellowish-green or golden. **Primary stems** short and inconspicuous, or conspicuous and creeping and spreading. **Secondary stems** erect to ascending, or stems and branches spreading or subascending, spirally foliate or complanate; pseudoparaphyllia apparently absent; radiculose below. **Leaves** ovate- to oblong-lanceolate or obovate-oblong symmetric or asymmetric, apex acuminate, acute or obtuse-apiculate, base slightly decurrent on one or both sides; margins plane to recurved, entire to bluntly or sharply serrate, bordered; costae single, 1/3-3/4 lamina length, sometimes forked along costa; laminal cells either short to ± long hexagonal and walls weak or lax, or cells oval to rhomboidal and walls firm; alar region undifferentiated (occasionally border cells at base numerous and extend part or fully across to costa). **Propagula** absent or present in leaf axial, short to long cylindrical. **Autoicous**, rarely dioicous. **Perichaetia** lateral; leaves differentiated, usually smaller than stem leaves. **Setae** elongate, slender to rather stout, smooth, papillose or ciliate distally. **Capsules** exserted, erect to pendulous, urn ovoid, neck distinct or not; exothecial cells usually collenchymatous; stomata present, at base of urn or on neck; annuli usually persistent, cells little differentiated; opercula conic-rostrate; peristome double, exostome teeth 16, papillose or striate and furrowed; endostome basal membrane low or high, segments 16, cilia reduced or absent. **Calyptrae** mitrate or campanulate, smooth or sparsely hairy, base fringed with hairs. **Spores** lightly to densely papillose, rarely smooth.

Plantas pequeñas hasta ocasionalmente medianas, formando manojos o matas delgadas, verde brillantes hasta verde-amarillentas o doradas. **Tallos primarios** cortos e inconspicuos, o conspicuos, reptantes y patentes. **Tallos secundarios** erectos hasta ascendentes, o tallos y ramas patentes o subascendentes, espiralmente foliados o complanados; pseudoparáfilos aparentemente ausentes; radiculosos hacia la base. **Hojas** ovado- hasta oblongo-lanceoladas u obovado-oblongas simétricas o asimétricas, ápices acuminados, agudos u obtuso-apiculados, bases escasamente decurrentes sobre uno o ambos lados; márgenes planos hasta recurvados, enteros hasta obtusamente o claramente serrados, bordeados; costa simple, ocupando 1/3-3/4 de la longitud de la lámina, algunas veces completamente bifurcadas; células de la lámina corto- hasta ± largo-hexagonales y con paredes débiles o laxas, o células ovales hasta rómbicas y con paredes firmes; región alar indiferenciada (ocasionalmente bordeada por numerosas células en la base y que se extienden parcial o totalmente hasta la costa). **Propágulos** ausentes o presentes en el eje axial o en las axilas de las hojas, corto- hasta largo-cilíndricos. **Autoicas**, raras veces dioicas. **Periquecios** laterales; hojas diferenciadas, generalmente menores que las hojas del tallo. **Setas** alargadas, delgadas hasta bastante robustas, lisas, distalmente papilosas o ciliadas. **Cápsulas** excertas, erectas hasta péndulas, urna ovoide, cuello conspicuo o no; células exoteciales generalmente colenquimatosas; estoma presentes, en la base de urna o sobre el cuello; anillo generalmente persistente, células poco diferenciadas; opérculo cónico-rostrado; peristoma doble, exostoma con 16 dientes, papilosos o estriados y acanalados; membrana basal del endostoma baja o alta, con 16 segmentos, cilios reducidos o ausentes. **Caliptra** mitrada o campanulada, lisa o con pelos esparcidos, base bordeada de pelos. **Esporas** finamente hasta densamente papilosas, rara vez lisas.