2. ***Pseudoleskeopsis zippelii*** (Dozy & Molk.) Broth. ni cao xian

Nat. Pflanzenfam. I(3): 1003. 1907. *Hypnum zippelii* Dozy & Molk., Ann. Sci. Nat. Bot., ser. 3, 2: 310. 1844. Type. Indonesia: Amboina, *Zippelinus s.n*.

Pseudoleskeopsis decurvata (Mitt.) Broth., Nat. Pflanzenfam. I(3): 1003. 1907. *Leskea decurvata* Mitt., J. Proc. Linn. Soc., Bot. 8: 154. 1864.

Pseudoleskeopsis integrifolia Broth., Symb. Sin. 4: 95. 1929. Type. China: Hunan,Yin-schan Wukang, *Handel-Mazzetti 12203* (holotype H).

Pseudoleskeopsis japonica (Sull.) Iwats., J. Hattori Bot. Lab. 29: 58. 1966. *Hypnum japonicum* Sull. in Perry, Narr. Exp. China Japan 2: 330. 1857.

Pseudoleskeopsis orbiculata (Mitt.) Broth., Nat. Pflanzenfam. I(3): 1003. 1907. *Hypnum orbiculatum* Mitt., J. Proc. Linn. Soc., Bot., Suppl. 1: 84. 1859.

Pseudoleskeopsis orbiculata var. *laticuspis* (Card.) Thér., Ann. Crypt. Exot. 2: 15. 1929. *Pseudoleskeopsis laticuspis* (Card.) Broth., Nat. Pflanzenfam. I(3): 1003. 1907. *Pseudoleskea laticuspis* Card., Beih. Bot. Centralbl. 17(1): 30. 1904.

Pseudoleskeopsis serrulata Card. & Thér., Bull. Acad. Int. Géogr. Bot. 21: 271. 1911. Type. China: Guizhou, Pingba, Chuts-shan, 10. 1. 1903, *Aortunat s.n*. (isotype H).

Plants rigid, dark green, yellowish green when young, in mats. *Stems* creeping, up to 3–4 cm long, irregularly branched; branches usually fascicular, erect, 0.5–1.0 cm long, slightly declinate when dry. *Stem leaves* loosely appressed on stems when dry, spreading when moist, triangularly ovate, 1.2–1.5 mm × 0.5–0.6 mm; upper margins crenate; costa thick, strong, ending below apex; branch leaves broadly ovate, involute or slightly secund when dry, spreading when moist, 1.0–1.4 mm × 0.5–0.8 mm, obtuse at apex; margins plane, usually dentate; costa ending before apex; upper and median leaf cells small, round hexagonal or rhomboid, 7–12 µm × 5–6 µm; basal leaf cells near margins subquadrate, along costa shortly rectangular. *Inner perichaetial leaves* lanceolate, 2–3 mm × 0.8 mm; cells linear, costa percurrent, broader below. *Setae* long 1.5–2.0 cm. reddish brown, sometimes two mature sporophytes borne in one perichaetium; capsules pendent, obovoid to oblong-ovoid, 1.5–2.0 mm long, with short, distinct apophysis, mouth contracted when dry; exostome and endostome equal in length, 0.4–0.6 mm long; exostome teeth lanceolate, nearly smooth, with lamella; endostome segments folded; basal membrane high; cilia 2. *Spores* spherical, 20–25 µm in diameter, verrucose.